

AD 741212

OFFICE OF NAVAL RESEARCH

Contract N00014-68-C-0250


Task No. NR 136-754

Site of Action of Plague Murine Toxin

by

S. Kadis

University of Georgia\*  
Department of Medical Microbiology  
College of Veterinary Medicine  
Athens, Georgia 30601



11 April 1972

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release; its  
distribution is unlimited

Reproduced by  
**NATIONAL TECHNICAL  
INFORMATION SERVICE**  
Springfield, Va. 22151

\*This work was completed at the Albert Einstein Medical Center  
in Philadelphia, Pennsylvania.

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Albert Einstein Medical Center York and Tabor Roads Philadelphia, Pennsylvania 19141		2a. REPORT SECURITY CLASSIFICATION No "restricted data" included	
3. REPORT TITLE  SITE OF ACTION OF PLAGUE MURINE TOXIN		2b. GROUP Information not available	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Final report - October 1967 - December 31, 1971			
5. AUTHOR(S) (First name, middle initial, last name)  Solomon Kadis			
6. REPORT DATE 11 April 1972		7a. TOTAL NO. OF PAGES 7	7b. NO. OF REFS 16
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)  7	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)  N/A	
10. DISTRIBUTION STATEMENT  This document has been approved for public release; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES  N/A		12. SPONSORING/MILITARY ACTIVITY Department of the Navy Office of Naval Research Arlington, Virginia 22217	
13. ABSTRACT Investigations on the mechanism of action of plague murine toxin have revealed that the toxin is capable of inhibiting the respiration of mitochondria isolated from toxin susceptible animals by complexing with a non-enzymatic protein, reported to be located in the membranes of mitochondria and to satisfy various criteria for a structural protein, and thereby altering the structural configuration of the multi-enzymatic unit of a mitochondrion. Studies on the effect of plague murine toxin on protein biosynthesis have shown that Toxin B, one of the two protein components of the plague toxin, inhibits the incorporation of $C^{14}$ leucine or $C^{14}$ phenylalanine in the presence of poly U into trichloroacetic acid precipitable material by cell-free rat liver systems. This inhibitory effect can be observed with as little as 10ug of toxin. The longer Toxin B is preincubated with the supernatant fraction, the greater is the percentage inhibition. This is not the case when Toxin B is preincubated with rat liver ribosomes. The inhibitory effect is not due to a protease or a nuclease activity of the toxin. In order to be able to trace the fate of plague murine toxin after injection to toxin susceptible mice and rats, an attempt was made to label the toxin with a gamma emitting isotope such as $Se^{75}$ -methionine. Examination of disc gel electrophoretic profiles of toxins A and B following the labeling procedures showed that in both cases one single protein could be seen and that practically all of the radioactivity was associated with these proteins. Both preparations retained high biological activity.			

The investigations which have been supported by this contract have been concerned with the mechanism of action of 2 protein toxins produced by the bacterium Pasteurella pestis, the causative agent of the disease known as plague. Our earlier studies have indicated that the toxin alters a number of mitochondrial energized processes primarily in heart mitochondria isolated from toxin-susceptible animals, but not in heart mitochondria isolated from toxin-resistant animals. The effects observed include inhibition of mitochondrial respiration (1, 2, 3), promotion of large amplitude mitochondrial swelling (4) and inhibition of respiration-dependent ion accumulation by mitochondria (5).

#### I. Interaction of Plague Murine Toxin with Mitochondrial Structural Protein

We have shown that the inhibition of mitochondrial respiration by plague murine toxin is related to its ability to alter the structural configuration of the multi-enzymatic unit of a mitochondrion. It was demonstrated that the toxin is able to interact with a non-enzymatic protein reported to be located in the membranes of mitochondria and to satisfy various criteria for a structural protein (6). This interaction with structural proteins derived from bovine or rat heart mitochondria results in the loss of toxicity of the toxin as well as the specific elimination of the gel diffusion precipitation band exhibited by Toxin B, one of the two protein species comprising plague murine toxin. Ultracentrifugal analysis of solutions containing equimolar ratios of Toxin B and bovine heart mitochondrial structural protein dissolved in 0.1% sodium dodecyl sulfate, 0.05 M NaCl and 0.01 M phosphate buffer (pH 10.5) revealed that the original 6.6 sedimentation coefficient (S) peak of the toxin and the 2.3 S peak of the structural protein vanished and a new 4.8 S peak was observed (7). The same results were obtained when the molar ratio of structural to toxin was 2:1. These findings suggest that structural protein is capable of complexing with Toxin B.

#### II. Effect of Plague Murine Toxin on the Incorporation of Amino Acids into Polypeptides by Cell-Free Rat Liver Systems

Reports on studies of the mechanism of action of bacterial protein toxins other than plague murine toxin have in recent years increasingly implicated some of these toxins in alterations involving protein biosynthesis by cells of certain tissues. During the past few years we have initiated experiments to determine if plague murine toxin has any effect on protein biosynthesis.

Initial amino acid incorporation studies were conducted with microsome and dialyzed 105,000 xg supernatant fluid utilizing a standard incorporation mixture similar to that described by Johnson et al. (8). With this system incorporation of  $C^{14}$  amino

acids into hot trichloroacetic acid (TCA)-precipitable materials is linear for approximately 20 minutes at 37°C, and is essentially complete after 40 minutes. The incorporation of C<sup>14</sup> amino acids was found to be highly sensitive to RNAase, puromycin and cycloheximide but virtually unaffected by large concentrations of purified DNAase.

Investigations on the effect of varying concentrations of Toxin B on the incorporation of C<sup>14</sup> leucine showed that when 10, 25, and 50 µg of toxin were incubated with 105,000 xg supernatant fluid for 20 minutes at 37°C prior to the initiation of the reaction, the rate of C<sup>14</sup> leucine incorporation was inhibited 30, 60 and 75%, respectively. Toxoid, formalin inactivated toxin, exhibits some inhibitory effect but approximately 10 to 20 times as much toxoid is needed to give an effect comparable to that of the active toxin. When C<sup>14</sup> phenylalanine incorporation by our cell-free rat liver system was measured in the presence of the artificial messenger, polyuridylic acid (poly U), the total incorporation in the absence of toxin at each time interval was appreciably higher than was observed in the C<sup>14</sup> leucine experiments. However, the percentage inhibition of incorporation in the presence of each concentration of toxin employed was essentially the same in both cases.

Experiments designed to determine whether or not the toxin has any effect on the reagents included in the reaction mixture such as ATP or the ATP-generating system rather than on protein biosynthesis per se were conducted by incubating all of the components of the reaction mixture exclusive of microsomes, supernatant fluid and C<sup>14</sup> amino acid with or without toxin for varying periods of time at 37°C and then allowing incorporation to take place by adding back the deleted constituents of the system. It was found that C<sup>14</sup> leucine was incorporated just as well in the presence of toxin as in its absence, indicating that the inhibitory effect of the toxin on amino acid incorporation into polypeptides is probably not due to any interaction of the toxin with any of the components of the incorporation mixture other than supernatant fluid or microsomes. Attempts were made to determine whether the time of preincubation of toxin with supernatant fluid is related to the extent of inhibition by the toxin of amino acid incorporation into polypeptides. In the absence of toxin, the incorporation of C<sup>14</sup> leucine does not vary with the time of preincubation of supernatant fluid at 37°C. In the presence of toxin, however, the inhibitory effect of a given concentration of toxin on C<sup>14</sup> leucine incorporation becomes more pronounced as the time of preincubation of the toxin with supernatant fluid is extended.

Experiments involving preincubation of the toxin with microsomes showed that although some inhibitory effect of the toxin could be noted, this did not vary with time of preincubation. The possibility exists that some supernatant fluid could have been associated with the microsomal pellets.

Since our cell-free rat liver protein synthesizing system is extremely sensitive to inhibition by very low concentrations of RNAase and RNAase is a frequent contaminant of many bacterial enzyme preparations, our Toxin B preparations were tested for RNAase activity by the procedure of Spahr and Schlessinger (9). After incubation of Toxin B alone, undialyzed 105,000 xg supernatant fluid alone or Toxin B together with supernatant fluid with  $C^{14}$  poly U and precipitation of the poly U with 60% cold ethanol, any  $C^{14}$  nucleotides or small oligonucleotides liberated by RNAase action were assayed as soluble radioactivity. The counts per minute (cpm) released following incubation of 145  $\mu$ g of Toxin B with  $C^{14}$  poly U were almost identical to that of the reagent blank. While undialyzed 105,000 xg supernatant fluid exhibits considerable RNAase activity, the addition of 145  $\mu$ g of Toxin B to this system did not increase the ethanol-soluble nucleotides liberated from  $C^{14}$  poly U. These results show that our Toxin B preparations are free from significant RNAase activity in addition to any factor that would liberate a latent RNAase in the supernatant fluid.

In order to be able to perform meaningful experiments involving the location of the site of inhibition by plague murine toxin of protein biosynthesis, cell-free systems from rat liver were assembled from better defined components in the particulate and soluble fractions. Amino acid incorporation studies were conducted with ribosomes purified essentially by the procedure of Weinstein (10) except that in the final stage the ribosomes were sedimented through a cushion of 10% sucrose in Tris buffer to which ammonium chloride, magnesium acetate and  $\beta$ -mercaptoethanol were added according to the procedure of Jones et al. (11). pH5 fractions were prepared by a modification of the procedure of Keller and Zamecnik (12) as described by Weinstein (10). Initial incorporation studies in the absence of toxin utilizing the incorporation mixture described by Weinstein (10) and the filter paper disc method of Mans and Novelli (13) for assaying radioactivity revealed that the above described purification procedures resulted in the removal of messenger RNA and transfer RNA (tRNA). Upon the addition of commercially available poly U and tRNA to a system containing the optimal ratio of ribosomes to pH5 fraction the incorporation of  $C^{14}$  phenylalanine into TCA precipitable material was appreciable and linear for approximately 60 minutes.

In the presence of 10, 25, and 50  $\mu$ g of Toxin B the percentage inhibition of  $C^{14}$  phenylalanine incorporation amounted to 20, 60 and 85%, respectively. Toxoid, on the other hand, had essentially no effect since as much as 200  $\mu$ g of the inactivated toxin inhibited amino acid incorporation by only 5 to 10%. These experiments involved the direct addition of the toxin or toxoid to the reaction mixture rather than preincubation with the pH5 fraction or the ribosomes.

### III. Distribution of Plague Murine Toxins In Vivo.

While there have been numerous studies conducted concerning the physiology and mechanism of action of plague toxins, the target organ or organs involved during infection with the plague bacillus or following toxin injection is still unknown. It is of interest, therefore, to prepare pure radioactively labeled toxin and trace its fate after injection to toxin susceptible mice and rats. It was envisaged that the detection procedure could be simplified by labeling the toxin with a gamma emitting isotope such as  $\text{Se}^{75}$ -methionine. We have carried out initial experiments (14) involving the labeling of Toxin A and Toxin B with  $\text{Se}^{75}$ -methionine which was included in the growth medium (15). The toxic proteins A and B were isolated by electrophoresis on polyacrylamide gel slabs and analyzed by disc gel electrophoresis (16). After the gels were stained, they were scanned with a Gilford spectrophotometer equipped with a linear transport attachment. Examination of the electrophoretic profiles of toxins A and B revealed that in both cases one single protein could be seen and that practically all of the radioactivity was associated with these proteins. The specific radioactivity was calculated to be 300cpm/g for Toxin A and 227 cpm/g for Toxin B. Both preparations retained high biological activity.

# REFERENCES CITED

1. Kadis, S., Ajl, S.J. and Rust, J.H., Jr. J. Bacteriol. 86, 757 (1963).
2. Kadis, S., Cohen, M. and Ajl, S. J. Biochim. Biophys. Acta 96, 179 (1965).
3. Kadis, S. and Ajl, S.J. J. Biol. Chem. 241, 1556 (1966).
4. Kadis, S. and Ajl, S.J. J. Biol. Chem. 238, 3472 (1963).
5. Kadis, S. Trenchard, A. and Ajl, S.J. Arch. Biochem. Biophys. 109, 272 (1965).
6. Kadis, S., Trenchard, A.V. and Ajl, S.J. J. Biol. Chem. 241, 5605 (1966).
7. Kadis, S., Urbano, C. and Ajl, S.J. Bacteriol. Proc. p. 88 (1967).
8. Johnson, W., Kuchler, R.J. and Solotorovsky, M. J. Bacteriol. 96, 1089 (1968).
9. Spahr, P.F. and Schlessinger, D. J. Biol. Chem. 238, PC 2251 (1963).
10. Weinstein, I.B. In "Methods in Enzymology" Vol. 12B (L. Grossman and K. Moldave, eds.) pp. 782-787, Academic Press, New York, N.Y.
11. Jones, O.W., Dieckmann, M. and Berg, P. J. Mol. Biol. 31, 177 (1968).
12. Keller, E.F. and Zamecnik, P.C. J. Biol. Chem. 221, 45 (1956).
13. Mans, R.J. and Novelli, G.D. Arch. Biochem. Biophys. 94, 48 (1961).
14. Leon, S.A., Kadis, S. and Kollmann, G. Nature 229, 120 (1971).
15. Englesberg, E. and Levy, J.B. J. Bacteriol. 68, 57 (1954).
16. Davis, B.J. Ann. N.Y. Acad. Sci. 121, 404 (1964).

PUBLICATIONS ISSUED UNDER CONTRACT

Kadis, S. and Ajl, S.J.

Alteration of mitochondrial respiration by plague murine toxin.  
J. Albert Einstein Medical Center, 15, 15-21 (1967).

Kadis, S., Urbano, C. and Ajl, S.J. (Abstract)

Ultracentrifugal analysis of complex formation between plague murine toxin and bovine heart mitochondrial structural protein.  
Bacteriol. Proc., p. 88 (1967).

Kadis, S. and Ajl, S.J. (Abstract)

Localization of the site of inhibition of heart mitochondrial respiration by plague murine toxin.  
VII International Congress for Biochemistry, Tokyo, Japan, Abstracts V, General sessions H-41, p. 89, (1967).

Kadis, S., Montie, T.C. and Ajl, S.J.

Plague Toxin.  
Scientific American, 220, 92-100 (1969).

Kadis, S. and Ajl, S.J. (Abstract)

Site of action of plague murine toxin.  
International Symposium on the Chemical Control of the Human Environment, Johannesburg, South Africa (1969).

Kadis, S. and Ajl, S.J.

Site and mode of action of murine toxin of Pasteurella pestis.  
In "Microbial Toxins" Volume 3 (T.C. Montie, S. Kadis and S.J. Ajl, eds.) pp. 39-67, Academic Press, New York, N.Y. (1970).

Kadis, S. and Trenchard, A.V., (Abstract)

Effect of plague murine toxin on incorporation of amino acids into polypeptides by cell-free rat liver system.  
Bacteriol Proc., p. 89 (1970).

Ajl, S.J. and Kadis, S. (Abstract)

Mode of action of plague murine toxin.  
X International Congress for Microbiology, Mexico City, Mexico (1970).

Leon, S.A., Kadis, S. and Kollmann, G.

Preparation of biologically active plague murine toxin labeled with <sup>75</sup>Se-methionine. Nature 229, 120-121 (1971).